

A Pattern Speed in the Galaxy's OH/IR Stars

Victor P. Debattista¹, Ortwin Gerhard¹, and Maartje N. Sevenster²

¹ Astronomisches Institut, Universität Basel, Venusstrasse 7, CH-4102 Binningen, Switzerland

² RSAA/MAAAO, RSAA/MSSSO, Cotter Road, Weston ACT 2611, Australia

1 Introduction

The Milky Way Galaxy (MWG) contains both a bar and spirals. The pattern speed/rotation frequency, Ω_p , of these components have been measured with a variety of models. For the bar, models have found values $40 \lesssim \Omega_p \lesssim 60$ km s⁻¹ kpc⁻¹ (Binney et al. 1991; Fux 1999; Englmaier & Gerhard 1999; Weiner & Sellwood 1999; Dehnen 1999; Bissantz et al. 2002). The spiral arm Ω_p is even more uncertain, with values in the range $13.5 \lesssim \Omega_p \lesssim 27$ km s⁻¹ kpc⁻¹ reported (Morgan 1990; Amaral & Lépine 1997; Mishurov & Zenina 1999).

A model-independent method, based on the continuity equation, for measuring pattern speeds in external galaxies has been developed by Tremaine & Weinberg (1984). This method can be extended to the MWG (Kuijken & Tremaine 1991; Debattista et al. 2002). For discrete tracers in the MWG, the Tremaine-Weinberg (TW) method is contained in the following expression:

$$\begin{aligned} \Delta V &\equiv \Omega_p R_0 - V_{\text{LSR}} \equiv \frac{\mathcal{K}}{\mathcal{P}} - u_{\text{LSR}} \frac{\mathcal{S}}{\mathcal{P}} \\ &= \frac{\sum_i f(r_i) v_{r,i}}{\sum_i f(r_i) \sin l_i \cos b_i} - u_{\text{LSR}} \frac{\sum_i f(r_i) \cos l_i \cos b_i}{\sum_i f(r_i) \sin l_i \cos b_i} \end{aligned} \quad (1)$$

where R_0 is the Sun-MWG center distance, V_{LSR} is the tangential velocity of the local standard of rest (LSR), u_{LSR} is the radial velocity of the LSR, $f(r_i)$ is the observational detection probability (which need not be known), v_r is the heliocentric radial velocity of a discrete tracer, and (l, b) are its Galactic coordinates. Eqn. 1 assumes only one pattern speed and a low amplitude for any rapidly growing structure. Moreover, the tracer population needs to be sampled uniformly; one such survey is the ATCA/VLA OH 1612 MHz survey (Sevenster et al. 1997a,b & 2001), covering $|l| \leq 45^\circ$ and $|b| \leq 3^\circ$.

2 Pattern Speed of the OH/IR Population

We extracted from the ATCA/VLA OH 1612 MHz survey a sample of ~ 250 OH/IR stars which are relatively old (older than 0.8 Gyr) and bright (flux density greater than 0.16 Jy). These selection criteria give OH/IR stars between 4 and 10 kpc away from the Sun. We applied the TW analysis of Eqn. 1 to this sample, obtaining the results shown in Fig. 1 (Debattista et al. 2002). Note that the




Fig. 1. The TW analysis for the OH/IR stars for changing l_0 (the maximum $|l|$ in Eqn. 1). From left to right are $\mathcal{P}$, $\mathcal{K}$, the number of stars and the resulting $\mathcal{K}/\mathcal{P}$

value of $\mathcal{K}/\mathcal{P}$ has converged, within the errors, for $|l| > 30^\circ$. Simple tests with models show that a measurement of ΔV with a sample of this size should give an average accuracy of $\sim 17\%$ and always better than 40% , when the asymmetry signal is as large as the one we find. From re-sampling experiments, we find $\Delta V = 252 \pm 41 \text{ km s}^{-1}$. If we assume $V_{\text{LSR}}/R_0 = 220/8 \text{ km s}^{-1} \text{ kpc}^{-1}$ (from SgrA* motion, Backer et al. 1999; Reid et al. 1999 and Cepheid proper motions, Feast & Whitelock 1997) and $u_{\text{LSR}} = 0$ (from SgrA* HI absorption spectrum, Radhakrishnan et al. 1980), we obtain $\Omega_p = 59 \pm 5 \text{ km s}^{-1} \text{ kpc}^{-1}$. We estimate systematic error to be $\sim 10 \text{ km s}^{-1} \text{ kpc}^{-1}$.

The signal we found is concentrated close the plane ($|b| \leq 1^\circ$) and at large longitude, suggesting a spiral is responsible for it (possibly the Scutum arm). The high Ω_p suggests this spiral arm is driven by the bar. Alternatively, the non-axisymmetric structure involved is an inner ring (Sevenster & Kalnajs 2001).

References

1. Amaral L. H., Lépine J. R. D., 1997, MNRAS, 286, 885
2. Backer D. C., Sramek R. A. 1999, ApJ, 524, 805
3. Binney J., Gerhard O. E., Stark A. A., et al. 1991, MNRAS, 252, 210
4. Bissantz N., Englmaier P., Gerhard O. 2002, MNRAS, *submitted*
5. Debattista, V. P., Gerhard, O. & Sevenster, M. N. 2001, MNRAS, *in press*
6. Dehnen W. 1999, ApJ, 524, L35
7. Englmaier P., Gerhard O. E. 1999, MNRAS, 304, 512
8. Feast M., Whitelock P. 1997, MNRAS, 291, 683
9. Fux R. 1999, A&A, 345, 787
10. Kuijken K., Tremaine S. 1991, *Dynamics of Disc Galaxies*, ed. B. Sundelius (Göteborg: Sweden) pg. 71
11. Mishurov Y. N., Zenina I. A. 1999, A&A, 341, 81
12. Morgan S. 1990, PASP, 102, 102
13. Radhakrishnan V., Sarma V. N. G. 1980, MNRAS, 85, 249
14. Reid M. J., Readhead A. C. S., Vermeulen R. C., et al. 1999, ApJ, 524, 816
15. Sevenster M. N., Kalnajs A. 2001, AJ, 122, 885
16. Sevenster M. N., Chapman J. M., Habing H. J., et al. 1997a, A&AS, 122, 79
17. Sevenster M. N., Chapman J. M., Habing H. J., et al. 1997b, A&AS, 124, 509
18. Sevenster M. N., van Langevelde H., Chapman J. M., et al. 2001, A&A, 366, 481
19. Tremaine S., Weinberg M. D. 1984, ApJ, 282, L5
20. Weiner B. J., Sellwood J. A. 1999, ApJ, 524, 112